

(19) World Intellectual Property Organization
International Bureau




(43) International Publication Date
13 September 2001 (13.09.2001)

PCT

(10) International Publication Number
WO 01/66540 A1

- (51) International Patent Classification⁷: C07D 401/04, 401/14, A61K 31/44, A61P 35/00
- (21) International Application Number: PCT/GB01/00916
- (22) International Filing Date: 2 March 2001 (02.03.2001)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
0005357.9 6 March 2000 (06.03.2000) GB
- (71) Applicant (for all designated States except US):
SMITHKLINE BEECHAM P.L.C. [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): GAIBA, Alessandra [IT/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). TAKLE, Andrew, Kenneth [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB). WILSON, David, Matthew [GB/GB]; GlaxoSmithKline, New Frontiers Science Park South, Third Avenue, Harlow, Essex CM19 5AW (GB).
- (74) Agent: BLAKEY, Alison, Jane; Corporate Intellectual Property, SmithKline Beecham, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: IMIDAZOL-2-CARBOXAMIDE DERIVATIVES AS RAF KINASE INHIBITORS



(I)

(57) Abstract: Compounds of the formula (I) wherein X is O, CH₂, S or NH, or the moiety X-R¹ is hydrogen; V is CH or N; R¹ is hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aryl, arylC₁₋₆alkyl, heterocyclyl, heterocyclylC₁₋₆alkyl, heteroaryl, or heteroarylC₁₋₆alkyl any of which except hydrogen may be optionally substituted; R² and R³ independently represent hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aryl, arylC₁₋₆alkyl, heteroaryl, heteroarylC₁₋₆alkyl, heterocyclyl, or heterocyclylC₁₋₆alkyl any one of which except hydrogen may be optionally substituted, or R² and R³ together with the nitrogen atom to which they are attached form a 4- to 10- membered op-

tionally substituted monocyclic or bicyclic ring; Ar is an aryl or heteroaryl ring either of which may be optionally substituted; one of X₁ and X₂ is N and the other is NR⁴, wherein R⁴ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl; or pharmaceutically acceptable salts thereof; their use as inhibitors of Raf kinases and pharmaceutical compositions containing them.

IMIDAZOL-2-CARBOXAMIDE DERIVATIVES AS RAF KINASE INHIBITORS


This invention relates to novel compounds and their use as pharmaceuticals particularly as Raf kinase inhibitors for the treatment of neurotraumatic diseases.

5 Raf protein kinases are key components of signal transduction pathways by which specific extracellular stimuli elicit precise cellular responses in mammalian cells. Activated cell surface receptors activate ras/rap proteins at the inner aspect of the plasmamembrane which in turn recruit and activate Raf proteins. Activated Raf proteins phosphorylate and activate the intracellular protein kinases MEK1 and MEK2. In turn, 10 activated MEKs catalyse phosphorylation and activation of p42/p44 mitogen-activated protein kinase (MAPK). A variety of cytoplasmic and nuclear substrates of activated MAPK are known which directly or indirectly contribute to the cellular response to environmental change. Three distinct genes have been identified in mammals that encode Raf proteins; A-Raf, B-Raf and C-Raf (also known as Raf-1) and isoformic 15 variants that result from differential splicing of mRNA are known.

Inhibitors of Raf kinases have been suggested for use in disruption of tumor cell growth and hence in the treatment of cancers, e.g. histiocytic lymphoma, lung adenocarcinoma, small cell lung cancer and pancreatic and breast carcinoma; and also in the treatment and/or prophylaxis of disorders associated with neuronal degeneration 20 resulting from ischemic events, including cerebral ischemia after cardiac arrest, stroke and multi-infarct dementia and also after cerebral ischemic events such as those resulting from head injury, surgery and/or during childbirth.

We have now found a group of novel compounds that are inhibitors of Raf kinases, in particular inhibitors of B-Raf kinase.

25 According to the invention there is provided compounds of formula (I):



(I)

wherein

X is O, CH₂, S or NH, or the moiety X-R¹ is hydrogen;

V is CH or N;

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aryl, arylC₁₋₆alkyl, heterocyclyl,
 5 heterocyclylC₁₋₆alkyl, heteroaryl, or heteroarylC₁₋₆alkyl any of which, except
 hydrogen, may be optionally substituted;

R² and R³ independently represent hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aryl,
 arylC₁₋₆alkyl, heteroaryl, heteroarylC₁₋₆alkyl, heterocyclyl, or heterocyclylC₁₋₆alkyl
 any one of which, except hydrogen, may be optionally substituted, or R² and R³
 10 together with the nitrogen atom to which they are attached form a 4- to 10-
 membered optionally substituted monocyclic or bicyclic ring;

Ar is an aryl or heteroaryl ring, either of which may be optionally
 substituted;

one of X₁ and X₂ is N and the other is NR⁴, wherein R⁴ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl;
 15 or pharmaceutically acceptable salts thereof.

As used herein, the double bond indicated by the dotted lines of formula (I),
 represent the possible tautomeric ring forms of the compounds falling within the
 20 scope of this invention, the double bond being to the unsubstituted nitrogen atom.

Alkyl and alkenyl groups referred to herein, individually or as part of larger
 groups e.g. alkoxy, may be straight or branched groups containing up to six carbon
 atoms and are optionally substituted by one or more groups selected from the group
 consisting of aryl, heteroaryl, heterocyclyl, C₁₋₆alkoxy, C₁₋₆alkylthio, arylC₁₋₆alkoxy,
 25 arylC₁₋₆alkylthio, amino, mono- or di-C₁₋₆alkylamino, cycloalkyl, cycloalkenyl,
 carboxy and esters thereof, amide, sulphonamido, ureido, guanidino, C₁₋₆alkylguanidino,
 amidino, C₁₋₆alkylamidino, C₁₋₆acyloxy, azido, hydroxy,
 hydroxyimino and halogen. Preferably the optional substituent contains a
 solubilising group; suitable solubilising moieties will be apparent to those skilled in
 30 the art and include hydroxy and amine groups. Even more preferably the optional

substituent include heterocyclyl e.g. piperidinyl, morpholinyl or piperazinyl, amino, mono- or di-C₁₋₆alkylamino, and hydroxy or any combination thereof.

Cycloalkyl and cycloalkenyl groups referred to herein include groups having from three to seven ring carbon atoms and are optionally substituted as described
5 hereinabove for alkyl and alkenyl groups.

When used herein, the term "aryl" includes, unless otherwise defined, single and fused rings suitably containing from 4 to 7, preferably 5 or 6, ring atoms in each ring, which rings, may each be unsubstituted or substituted by, for example, up to three substituents.

10 Suitable aryl groups include phenyl and naphthyl such as 1-naphthyl or 2-naphthyl.

When used herein the term "heterocyclyl" includes, unless otherwise defined, non-aromatic, single and fused, rings suitably containing up to four heteroatoms in each ring, each of which is selected from O, N and S, which rings,
15 may be unsubstituted or substituted by, for example, up to three substituents. Each heterocyclic ring suitably has from 4 to 7, preferably 5 or 6, ring atoms. A fused heterocyclic ring system may include carbocyclic rings and need include only one heterocyclic ring. Examples of heterocyclyl groups include pyrrolidine, piperidine, piperazine, morpholine, imidazolidine and pyrazolidine.

20 When used herein, the term "heteroaryl" includes, unless otherwise defined, mono- and bicyclic heteroaromatic ring systems comprising up to four, preferably 1 or 2, heteroatoms each selected from O, N and S. Each ring may have from 4 to 7, preferably 5 or 6, ring atoms. A bicyclic heteroaromatic ring system may include a carbocyclic ring. Examples of heteroaryl groups include pyrrole, quinoline,
25 isoquinoline, pyridine, pyrimidine, oxazole, thiazole, thiadiazole, triazole, imidazole and benzimidazole.

Aryl, heterocyclyl and heteroaryl groups may be optionally substituted by preferably up to three substituents. Suitable substituents include halogen, C₁₋₆alkyl, aryl, aryl C₁₋₆alkyl, C₁₋₆alkoxy, C₁₋₆alkoxy C₁₋₆alkyl, halo C₁₋₆alkyl, arylC₁₋₆alkoxy,
30 hydroxy, nitro, cyano, azido, amino, mono- and di-N-C₁₋₆alkylamino, acylamino,

arylcarbonylamino, acyloxy, carboxy, carboxy salts, carboxy esters, carbamoyl,
 mono- and di-*N*-C₁₋₆alkylcarbamoyl, C₁₋₆alkoxycarbonyl, aryloxycarbonyl, ureido,
 guanidino, C₁₋₆alkylguanidino, amidino, C₁₋₆alkylamidino, sulphonylamino,
 aminosulphonyl, C₁₋₆alkylthio, C₁₋₆alkylsulphinyl, C₁₋₆alkylsulphonyl, heterocyclyl,
 5 heteroaryl, heterocyclyl C₁₋₆alkyl, hydroxyimino-C₁₋₆alkyl and heteroaryl C₁₋₆alkyl
 and combinations thereof. Preferably the optional substituent contains a solubilising
 group; suitable solubilising moieties will be apparent to those skilled in the art and
 include hydroxy and amine groups. Even more preferably the optional substituent
 include heterocyclyl, amino, mono- or di-C₁₋₆alkylamino, amide, and hydroxy or any
 10 combination thereof.

X is preferably NH or X-R¹ is preferably hydrogen and when X is NH, R¹ is
 preferably hydrogen or C₁₋₆alkyl.

When V is CH, X-R¹ is preferably hydrogen.

When V is N, X-R¹ is preferably NH₂.

15 Most preferably X-R¹ is hydrogen.

Ar is preferably an optionally substituted phenyl.

Preferred substituents for the group Ar include halo, hydroxy, hydroxy C₁₋₆
 alkyl e.g. hydroxymethyl, hydroxyimino-C₁₋₆alkyl and C₁₋₆alkoxy e.g. methoxy.

20 More preferred are halo and hydroxy. When Ar is phenyl the substituents are
 preferably present in the 3-position or the 3,4-positions. When Ar is phenyl it
 preferably has a 3-hydroxy or 3-chloro substituent, more preferably a 3-hydroxy
 substituent. Particular substitution patterns for Ar when phenyl are 3-hydroxy, 3-
 hydroxy-4-halo e.g. 3-hydroxy-4-chloro or 3-hydroxy-4-bromo, 3-hydroxy-4-methyl
 and 3-hydroxy-4-methoxy, more particularly 3-hydroxy-4-chloro.

25 Preferably R² and R³ independently represent hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl,
 aryl, arylC₁₋₆alkyl, heteroaryl, heteroarylC₁₋₆alkyl, heterocyclyl, or heterocyclylC₁₋₆alkyl, or
 R² and R³ together with the nitrogen atom to which they are attached form a 4- to 10-
 membered monocyclic or bicyclic ring.

30 Even more preferably R² and R³ independently represent hydrogen, C₁₋₆alkyl, aryl,
 heteroarylC₁₋₆alkyl, any of which except hydrogen can be optionally substituted or R² and

R³ together with the nitrogen atom to which they are attached form an optionally substituted 5 or 6 membered monocyclic or bicyclic ring for example piperidine.

The compounds of formula (I) preferably have a molecular weight of less than 800.

Particular compounds according to the invention include those mentioned in
5 the examples and their pharmaceutically acceptable salts. It will be understood that the invention includes pharmaceutically acceptable derivatives of compounds of formula (I) and that these are included within the scope of the invention.

As used herein "pharmaceutically acceptable derivative" includes any pharmaceutically acceptable salt, ester or salt of such ester of a compound of
10 formula (I) which, upon administration to the recipient, is capable of providing (directly or indirectly) a compound of formula (I) or an active metabolite or residue thereof.

It will be appreciated that for use in medicine the salts of the compounds of formula (I) should be pharmaceutically acceptable. Suitable pharmaceutically
15 acceptable salts will be apparent to those skilled in the art and include those described in *J. Pharm. Sci.*, 1977, 66, 1-19, such as acid addition salts formed with inorganic acids e.g. hydrochloric, hydrobromic, sulfuric, nitric or phosphoric acid; and organic acids e.g. succinic, maleic, acetic, fumaric, citric, tartaric, benzoic, p-toluenesulfonic, methanesulfonic or naphthalenesulfonic acid. Other salts e.g.
20 oxalates, may be used, for example in the isolation of compounds of formula (I) and are included within the scope of this invention.

The compounds of this invention may be in crystalline or non-crystalline form, and, if crystalline, may optionally be hydrated or solvated. This invention
includes within its scope stoichiometric hydrates as well as compounds containing
25 variable amounts of water.


The invention extends to all isomeric forms including stereoisomers and geometric isomers of the compounds of formula (I) including enantiomers and mixtures thereof e.g. racemates. The different isomeric forms may be separated or resolved one from the other by conventional methods, or any given isomer may be

obtained by conventional synthetic methods or by stereospecific or asymmetric syntheses.

Since the compounds of formula (I) are intended for use in pharmaceutical compositions it will readily be understood that they are each preferably provided in substantially pure form, for example at least 60% pure, more suitably at least 75%
5 pure and preferably at least 85%, especially at least 98% pure (% are on a weight for weight basis). Impure preparations of the compounds may be used for preparing the more pure forms used in the pharmaceutical compositions.

Compounds of formula (I) are imidazole derivatives which may be readily
10 prepared using procedures well-known to those skilled in the art, and described in, for instance, Comprehensive Heterocyclic Chemistry, Editors Katritzky and Rees, Pergamon Press, 1984, 5, 457-497, from starting materials which are either commercially available or can be prepared from such by analogy with well-known processes. A key step in many such syntheses is the formation of the central
15 imidazole nucleus. Suitable procedures are described in *inter alia* US Patent No's. 3,707,475 and 3,940,486 which are herein incorporated by reference in their entirety. These patents describe the synthesis of α -diketones and α -hydroxyketones (benzoins) and their subsequent use in preparing imidazoles and N-hydroxyl imidazoles.

20



Preferred methods for preparing compounds of this invention are as outlined in the above scheme, wherein R is C₁₋₆ alkyl or arylC₁₋₆ alkyl, preferably ethyl. α-Diketones are prepared by condensation of the anion of, for example, a 4-substituted pyridine derivative (V = CH, R¹-X = H) with the Weinreb amide of an aryl acid, or with an aryl-aldehyde, followed by oxidation of the intermediate product. Stirring the diketone with an aldehyde, such as glyoxylic acid ethyl ester, and ammonium acetate in a mixture of methanol and methyl-*tert*-butyl ether allows access to the imidazole nucleus, by analogy to the method described in patent WO 98/56788. Thereafter, the ethyl ester, the corresponding acid or an activated derivative thereof may be converted into an amide using conventional amide bond forming procedures. Such procedures are well known in the art and are described in, for instance, P.D. Bailey, I.D. Collier and K.M. Morgan in *Comprehensive Organic Functional Group Transformation*, Vol. 5, ed. C.J. Moody, p.257, Elsevier Scientific, Oxford, 1995.

Non-selective alkylation of the imidazole nitrogen (using one of the procedures outlined in N. J. Liverton et al; *J. Med. Chem.*, 1999, 42, 2180-2190) with a compound of formula L-R⁴ wherein L is a leaving group, e.g. halo, sulfonate or triflate, will yield both isomers of the compounds of formula (I) where X₁ or X₂ is

NR⁴ in which R⁴ is other than hydrogen, the isomers can be separated by chromatographic methods.


During the synthesis of the compounds of formula (I) labile functional groups in the intermediate compounds, e.g. hydroxy, carboxy and amino groups, may be protected. A comprehensive discussion of the ways in which various labile functional groups may be protected and methods for cleaving the resulting protected derivatives is given in for example *Protective Groups in Organic Chemistry*, T.W. Greene and P.G.M. Wuts, (Wiley-Interscience, New York, 2nd edition, 1991).

The compounds of formula (I) may be prepared singly or as compound libraries comprising at least 2, for example 5 to 1,000 compounds, and more preferably 10 to 100 compounds of formula (I). Libraries of compounds of formula (I) may be prepared by a combinatorial 'split and mix' approach or by multiple parallel synthesis using either solution phase or solid phase chemistry, by procedures known to those skilled in the art.

Thus according to a further aspect of the invention there is provided a compound library comprising at least 2 compounds of formula (I), or pharmaceutically acceptable salts thereof.

Pharmaceutically acceptable salts may be prepared conventionally by reaction with the appropriate acid or acid derivative.

The novel carboxylic esters and the corresponding acids of formula (II) which are used as intermediates in the synthesis of the compounds of formula (I) also form part of the present invention:



25

(II)

wherein X, V, R¹, Ar, X₁ and X₂ are as defined for formula (I) and R is hydrogen, C₁₋₆ alkyl or arylC₁₋₆ alkyl.

As indicated above the compounds of formula (I) and their pharmaceutically acceptable salts are useful for the treatment and/or prophylaxis of disorders in which Raf kinases, in particular B-Raf kinase, are implicated.

According to a further aspect of the invention there is provided the use of a
5 compound of formula (I) or a pharmaceutically acceptable salt thereof as an inhibitor of B-Raf kinase.

As indicated above the compounds of formula (I) and their pharmaceutically acceptable salts are useful the treatment and/or prophylaxis of disorders associated with neuronal degeneration resulting from ischemic events.

10 According to a further aspect of the invention there is provided a method of treatment or prophylaxis of a neurotraumatic disease, in a mammal in need thereof, which comprises administering to said mammal an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof.

According to a further aspect of the invention there is provided the use of a
15 compound of formula (I) or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by a neurotraumatic event.

Neurotraumatic diseases/events as defined herein include both open or penetrating head trauma, such as caused by surgery, or a closed head trauma injury, such
20 as caused by an injury to the head region. Also included within this definition is ischemic stroke, particularly to the brain area, transient ischemic attacks following coronary by-pass and cognitive decline following other transient ischemic conditions.

Ischemic stroke may be defined as a focal neurologic disorder that results from insufficient blood supply to a particular brain area, usually as a consequence of an
25 embolus, thrombi, or local atheromatous closure of the blood vessel. Roles for stress stimuli (such as anoxia), redox injury, excessive neuronal excitatory stimulation and inflammatory cytokines in this area has been emerging and the present invention provides a means for the potential treatment of these injuries. Relatively little treatment, for an acute injury such as these has been available.

The compounds of the invention may also be used in the treatment or prophylaxis of cancers.

The compounds of the invention may also be of use for the treatment or prophylaxis of CSBP/p38 mediated diseases as described in WO 99/01131 and WO
5 99/01130.

It will be appreciated by those skilled in the art that reference herein to treatment extends to prophylaxis as well as the treatment of established infections or symptoms.

In order to use the compounds of formula (I) in therapy, they will normally
10 be formulated into a pharmaceutical composition in accordance with standard pharmaceutical practice.

According to a further aspect of the invention there is provided a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.

15 The compounds of formula (I) may conveniently be administered by any of the routes conventionally used for drug administration, for instance, parenterally, orally, topically or by inhalation. The compounds of formula (I) may be administered in conventional dosage forms prepared by combining it with standard pharmaceutical carriers according to conventional procedures. The compounds of
20 formula (I) may also be administered in conventional dosages in combination with a known, second therapeutically active compound. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the pharmaceutically acceptable carrier is dictated by the amount of compound of
25 formula (I) with which it is to be combined, the route of administration and other well-known variables. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

The pharmaceutical carrier employed may be, for example, either a solid or
30 liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar,

pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax.

5 A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25mg to about 1g. When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile
10 injectable liquid such as an ampoule or nonaqueous liquid suspension.

 The compounds of formula (I) are preferably administered parenterally, that is by intravenous, intramuscular, subcutaneous intranasal, intrarectal, intravaginal or intraperitoneal administration. The intravenous form of parenteral administration is generally preferred. The compounds may be administered as a bolus or continuous
15 infusion e.g. over 3 days. Appropriate dosage forms for such administration may be prepared by conventional techniques.

 The compounds of formula (I) may also be administered orally. Appropriate dosage forms for such administration may be prepared by conventional techniques.

 The compounds of formula (I) may also be administered by inhalation, that is by
20 intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as aerosol formulations, may be prepared by conventional techniques.

 The compounds of formula (I) may also be administered topically, that is by non-systemic administration. This includes the application of the inhibitors externally to the
25 epidermis or the buccal cavity and the instillation of such a compound into the ear, eye and nose, such that the compound does not significantly enter the blood stream.

 For all methods of use disclosed herein the daily oral dosage regimen will preferably be from about 0.1 to about 80 mg/kg of total body weight, preferably from about 0.2 to 30 mg/kg, more preferably from about 0.5mg to 15mg. The daily parenteral
30 dosage regimen about 0.1 to about 80 mg/kg of total body weight, preferably from about

0.2 to about 30 mg/kg, and more preferably from about 0.5mg to 15mg/kg. The daily topical dosage regimen will preferably be from 0.1 mg to 150mg, administered one to four, preferably two or three times daily. The daily inhalation dosage regimen will preferably be from about 0.01 mg/kg to about 1 mg/kg per day. It will also be

5 recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of the inhibitors will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular patient being treated, and that such optimums can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the

10 number of doses of the inhibitors given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests. In the case of pharmaceutically acceptable salts the above figures are calculated as the parent compound of formula (I).

No toxicological effects are indicated/expected when a compound of formula

15 (I) is administered in the above mentioned dosage range.

All publications, including but not limited to patents and patent applications, cited in this specification are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

20 The following Examples illustrate the preparation of pharmacologically active compounds of the invention and the following Descriptions illustrate the preparation of intermediates used in the preparation of these compounds.

Abbreviations used herein are as follows;

25 THF means tetrahydrofuran.

Description 1: 5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1*H*-imidazole-2-carboxylic acid ethyl ester

Step 1. 1-(3,4-Dichlorophenyl)-2-pyridin-4-yl-ethane-1,2-diol

4-(*tert*-Butyldimethylsilyloxymethyl)-pyridine (T. F. Gallagher *et al*, *Bioorganic and Medicinal Chemistry*; 1997, 5, 49) (67g, 0.3mol) was dissolved in THF (250ml) and cooled to -40°C. The solution was treated with a 2M solution of lithium diisopropylamide in THF (200ml, 0.4mol) and stirred for 45 min maintaining a temperature of -40 to -20°C, before the dropwise addition of 3,4-dichlorobenzaldehyde (55.13g, 0.32mol) in THF (250ml). The mixture was allowed to warm to room temperature then stirred for a further 18 hours. After re-cooling to 0°C the reaction was quenched with saturated ammonium chloride solution (500ml), and the resulting two phase mixture separated. The aqueous phase was extracted with ethyl acetate and the combined organics concentrated under reduced pressure. The residue was dissolved in ethyl acetate, washed with saturated sodium bicarbonate solution, water and brine, dried (MgSO₄) and concentrated under reduced pressure to an oil (129g). The oil was dissolved in THF (300ml) and a 1M solution of tetrabutylammonium fluoride (360ml, 0.36mol) added dropwise. The solution was stirred at room temperature for 45 min, then concentrated to an oil under reduced pressure. The oil was dissolved in ethyl acetate and washed with saturated sodium bicarbonate solution, water and brine, dried (MgSO₄) and evaporated under reduced pressure. The oil was triturated with hexane and the resulting solid filtered and washed with hexane to afford the title compound (67.58g 79%) as a tan solid; MS(AP+) m/e 284/286/288 [M+H]⁺.

Step 2. 1-(3,4-Dichlorophenyl)-2-pyridin-4-yl-ethane-1,2-dione

Dimethylsulfoxide (37ml, 0.53mol) was dissolved in dichloromethane (250ml) and cooled to -78°C. Oxalyl chloride (34.5ml, 0.40mol) was added dropwise and the solution stirred for 20 min. A solution of the product of Step 1 (34g, 0.12mol) in dimethylsulfoxide (40ml) and dichloromethane (200ml) was added dropwise at -78°C, and the solution stirred for 30 min. Triethylamine (104ml, 0.74mol) was added dropwise and the solution became flocculent such that overhead stirring became necessary. The solution was allowed to stir at room temperature over 2 hours then was poured on to ice/saturated sodium bicarbonate solution. The aqueous layer was separated, and re-extracted with dichloromethane. The combined

organic phases were concentrated under reduced pressure to a green-yellow solid. The solid was redissolved in dichloromethane and washed with water and brine, dried (MgSO_4) and evaporated to a solid. The crude solid was purified by silica gel chromatography eluting with dichloromethane, to afford the title compound (28.6g, 5 85%) as a yellow solid; MS(-ve ion) m/e 279/281/283 $[\text{M-H}]^-$.

Step 3. 5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid ethyl ester

The product of Step 2 (2.0g, 7.1mmol) was dissolved in *tert*-butyl-methyl ether (50ml). Glyoxylic acid ethyl ester (50% solution in toluene, 2.8ml, 14.3mmol) was 10 then added followed by a solution of ammonium acetate (1.37g, 17.8mmol) in methanol (10ml) which was added dropwise over 30 min. After stirring at room temperature overnight, the solvent was evaporated in vacuo, the residue dissolved in dichloromethane and then washed with aqueous potassium carbonate and brine. The organic layer was dried (MgSO_4), filtered and concentrated at reduced pressure. The 15 residue was chromatographed on silica gel eluting with 5% of a 9:1 methanol : 0.880ammonia solution, in ether to give the title compound as a pale yellow solid (600mg, 23%); MS(AP+) m/e 362/364 $[\text{M+H}]^+$.

Description 2: 5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid 20

D1 (520mg, 1.4mmol) was dissolved in methanol (10ml), 40% aqueous sodium hydroxide solution (2ml) added and the mixture heated to reflux overnight. On cooling, the solvent was removed in vacuo, the residue dissolved in water and washed with ether. The aqueous layer was cooled to 0°C and acidified with acetic 25 acid. The precipitate was filtered and dried under vacuum overnight to give the title compound, as a white solid (400mg, 83%); MS(AP+) m/e 334 $[\text{M+H}]^+$.

Description 3: 5-(4-Chloro-3-methoxyphenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid ethyl ester

30 **Step 1. 4-Chloro-3-,N-dimethoxy-N-methyl-benzamide**

A suspension of 4-chloro-3-methoxybenzoic acid (F. Claudi *et al J. Med. Chem.*, 1992, 35, 4408) (37.2g, 0.2mol) in dichloromethane (500ml) containing oxalyl chloride (26ml) was treated with *N,N*-dimethylformamide (10 drops). After stirring at room temperature for 6 hours the solution was concentrated at reduced pressure, additional dichloromethane was added to the residue and the solvent was re-evaporated. The residue was then dissolved in acetonitrile (600ml) and methoxymethylamine hydrochloride (20.5g, 0.21mol) added. The mixture was cooled in an ice-bath, a solution of pyridine (80ml) in acetonitrile (150ml) added dropwise, and the mixture stirred at room temperature for 18 hours. The solution was concentrated and the residue partitioned between ethyl acetate and saturated potassium carbonate solution. The organic layer was separated, washed with brine, dried (MgSO_4) and concentrated at reduced pressure to give the title compound (40.0g, 87%) as a colourless oil; MS(ES+) m/e 230/232 $[\text{M}+\text{H}]^+$.

Step 2. 1-(4-Chloro-3-methoxy-phenyl)-2-pyridin-4-yl-ethanone

4-Picoline (16.9ml, 0.174mol) was added dropwise to a stirred solution of lithium di-isopropylamide (110ml, 0.22mol, 2M solution in heptane, ethylbenzene, tetrahydrofuran) in dry tetrahydrofuran (150ml) at -78°C . After stirring at -78°C for 15 min a solution of the product of Step 1 (40.0g, 0.174mol) in tetrahydrofuran (100ml) was added dropwise. The reaction was allowed to warm to room temperature over 3 hours. The solution was cooled in ice and saturated ammonium chloride solution added. The aqueous mixture was extracted with ethyl acetate, washed with brine, dried (MgSO_4), filtered and concentrated at reduced pressure. The resulting gum was triturated with cold diethyl ether/hexane (1:1, 300ml) and the solid collected to give the title compound, as a pale yellow solid (29g, 64%); MS(ES+) m/e 262/264 $[\text{M}+\text{H}]^+$.

Step 3. 1-(4-Chloro-3-methoxy-phenyl)-2-pyridin-4-yl)-ethane-1,2-dione

A solution of the product of Step 2 (22.5g, 86mmol) in dimethylsulphoxide (150ml) was stirred at 55°C . Hydrogen bromide (48% aqueous, 21ml) was added dropwise and the solution maintained at 55°C for 1 hour. After cooling to room temperature, the solution was poured into a solution of sodium acetate (21g) in ice-water (1 litre)

and the resulting slurry was stirred at room temperature for 30 min. The mixture was extracted with ethyl acetate and the organic layers were combined, washed with brine, dried (MgSO₄), filtered and concentrated at reduced pressure. The residue was triturated with diethyl ether/hexane (1:4) and the solid collected to give the title
5 compound as a yellow solid; MS(EI) m/e 275/277 [M]⁺.

Step 4. 5-(4-Chloro-3-methoxyphenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid ethyl ester

The title compound (120mg, 37%) was prepared from the product of Step 3 and glyoxylic acid ethyl ester using the method described in D1 Step 3; MS(ES⁺) m/e
10 358/360/362 [M+H]⁺.

Description 4: 5-(4-Chloro-3-methoxyphenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid

The title compound (120mg, 78%) was prepared from the product of D3 and
15 aqueous sodium hydroxide, using the method described in D2; MS(AP⁻) m/e 328 [M-H]⁻.

Example 1: 1-[5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazol-2-yl]-1-piperidin-1-yl-methanone

20 **Step 1. 5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazole-2-carbonyl chloride**

D2 (360mg, 1.1mmol) was dissolved in dichloromethane (15ml), oxalyl chloride (0.3ml, 3.3mmol) was added, followed by 1 drop of DMF, and the mixture stirred at room temperature for 1h. The solvent was evaporated to give the title compound as
25 a yellow solid which was used directly in the next step.

Step 2. 1-[5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazol-2-yl]-1-piperidin-1-yl-methanone

The product of Step 1 (52mg, 0.15mmol) was dissolved in anhydrous dichloromethane (3ml), piperidine (0.025ml, 0.23mmol) was added followed by
30 triethylamine (0.084ml, 0.6mmol) and the mixture stirred at room temperature for 2

hours. The mixture was diluted with dichloromethane and washed with aqueous sodium bicarbonate and brine. The organic layer was dried (MgSO_4) and evaporated under vacuum. The residue was chromatographed on silica gel eluting with 5% of a 9:1 methanol:0.880 ammonia solution, in dichloromethane to give the title
5 compound as a white solid (16mg, 26%); MS(AP+) m/e 402 $[\text{M}+\text{H}]^+$.

Example 2: 5-(4-Chloro-3-methoxyphenyl)-4-pyridin-4-yl-1*H*-imidazol-2-yl]-1-piperidin-1-yl-methanone

Step 1. 5-(4-Chloro-3-methoxyphenyl)-4-pyridin-4-yl-1*H*-imidazole-2-
10 **carbonyl chloride**

The title compound was prepared from the product of D4 using the method described in Example 1 Step 1.

Step 2. 5-(4-Chloro-3-methoxyphenyl)-4-pyridin-4-yl-1*H*-imidazol-2-yl]-1-piperidin-1-yl-methanone

15 The title compound (30mg, 50%) was prepared from the product of Step 1 and piperidine using the method described in Example 1, Step 2: MS(AP+) m/e 397/399 $[\text{M}+\text{H}]^+$.

Example 3: 1-[5-(4-Chloro-3-hydroxyphenyl)-4-pyridin-4-yl-1*H*-imidazol-2-
20 **yl]-1-piperidin-1-yl-methanone**

A solution of Example 2 (18mg, 0.045mmol) in dichloromethane (5ml) was cooled to 0°C and treated with boron tribromide (1M in dichloromethane, 0.14ml, 0.14mmol). The solution was stirred at 0°C for 1 hour then at room temperature for a further 5 hours. 2M Hydrochloric acid (1ml) was added and the reaction heated to
25 50°C for 1 hour. After cooling, the mixture was basified with potassium carbonate solution and the resultant precipitate collected by filtration. The residue was chromatographed on silica gel eluting with 10% of a 9:1 methanol:0.880 ammonia solution, in dichloromethane to give the title compound as a white solid (10mg, 58%); MS(AP+) m/e 383/385 $[\text{M}+\text{H}]^+$.

30

The following examples were prepared by the general two-step method described in Example 1.

Example	Amine	Characterisation
4	5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid (3-dimethylamino-propyl)-amide	3-Dimethyl propylamine MS(AP+) m/e 418/420 [M+H] ⁺
5	3-({1-[5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazol-2-yl]-methanoyl}-amino)-propionic acid tert-butyl ester	3-Amino-propionic acid tert-butyl ester MS(AP+) m/e 462/464 [M+H] ⁺
6	3-({1-[5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazol-2-yl]-methanoyl}-amino)-propionic acid	4-Amino-propionic acid MS(AP+) m/e 405/407 [M+H] ⁺
7	5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid [2-(1H-indol-3-yl)-ethyl]-amide	2-(1H-Indol-3-yl)-ethylamine MS(AP+) m/e 476/478 [M+H] ⁺
8	5-(3,4-Dichlorophenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid (4-morpholin-4-yl-phenyl)-amide	4-Morpholin-4-yl-phenylamine MS(AP+) m/e 497 [M+H] ⁺

5 The following examples were prepared by the general two-step method described in Example 2.

Example	Amine	Characterisation
9	5-(4-Chloro-3-methoxyphenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid (3-dimethylamino-propyl)-amide	3-Dimethyl aminopropyl amine MS(AP+) m/e 415/417 [M+H] ⁺

The following examples were prepared by the general method described in Example 3.

Example	Precursor	Characterisation
10 5-(4-Chloro-3-hydroxyphenyl)-4-pyridin-4-yl-1 <i>H</i> -imidazole-2-carboxylic acid (3-dimethylamino-propyl)-amide	Example 9	MS(AP+) m/e 400/402 [M+H] ⁺

5 **Example 11: 5-(4-chloro-3-hydroxyphenyl)-4-pyridin-4-yl-1*H*-imidazole-2-carboxylic acid (2-dimethylamino ethyl) amide**

Step 1. 5-(4-Chloro-3-hydroxyphenyl)-4-pyridin-4-yl-1*H*-imidazole-2-carboxylic acid.

- 10 D3 (7.60g, 21 mmol) was suspended in dichloromethane (200 ml) and treated with a 1 molar solution of boron tribromide in dichloromethane (100 ml, 100 mmol). The suspension was stirred at room temperature for 16 hours, 5M hydrochloric acid (50 ml) added and the mixture heated at reflux for a further 30 minutes. The mixture was then concentrated and the pH adjusted to ca. pH11 with 40% sodium hydroxide
- 15 solution. The solution was warmed to 50°C to complete ester hydrolysis and the product precipitated by acidification to pH4-5 with acetic acid. The solid was filtered, washed with water and ether then dried under vacuum to afford the title compound (3.67g, 55%); MS(AP+) m/e 315/317 [M+H]⁺

20 **Step 2. 5-(4-Chloro-3-hydroxyphenyl)-4-pyridin-4-yl-1*H*-imidazole-2-carbonyl chloride**

- A suspension of the product from Step 1 (1.24g, 3.9mmol) in dichloromethane (50ml) was treated with oxalyl chloride (3.4ml, 39mmol) and 10 drops dimethylformamide. The suspension was then heated at reflux for 5 hours, cooled
- 25 and concentrated under vacuum. The resultant residue was azeotroped with toluene

then triturated with hexane to afford the title compound (1.56g), which was used without further purification.

Step 3. 5-(4-chloro-3-hydroxyphenyl)-4-pyridin-4-yl-1H-imidazole-2-

5 carboxylic acid (2-dimethylamino ethyl) amide

The product from Step 2 (124 mg, 0.3 mmol) was suspended in dichloromethane (5 ml) and treated with a solution of N,N-dimethyl ethylenediamine (0.036 ml, 0.33 mmol), and triethylamine (0.125ml, 0.9mmol) in dichloromethane (1ml). The reaction mixture was then stirred at room temperature overnight, washed saturated sodium bicarbonate solution (3 ml) and passed through a Varain Chem Elut hydromatric cartridge to remove the aqueous phase. The filtrate was then concentrated and purified by silica gel chromatography, eluting with 0.3:3:7 dichloromethane: ethanol:0.880 ammonia solution to afford the title compound (20mg, 17%); MS(AP+) m/e 386/388 [M+H]⁺

The following examples were prepared by the general method described in Example 11.

Example		Amine	Characterisation
12	5-(4-chloro-3-hydroxy-phenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid (3-piperidin-1-yl-propyl) amide	3-(piperidin-1-yl)-propylamine	MS(AP+) m/e 440/442 [M+H] ⁺
13	5-(4-chloro-3-hydroxyphenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid (3-dimethylamino-2-dimethylpropyl) amide	N,N,-2,2-tetramethyl-propane-1,3-diamine	MS(AP+) m/e 428/430 [M+H] ⁺
14	5-(4-chloro-3-hydroxyphenyl)-4-pyridin-4-yl-1H-imidazole-2-carboxylic acid (4-benzyl-piperazin-1-yl) amide	N-benzyl piperazine	MS(AP+) m/e 474/476 [M+H] ⁺

It is to be understood that the present inventions covers all combinations of particular and preferred subgroups described hereinabove.

BIOLOGICAL EXAMPLES

- 5 The activity of compounds of formula (I) as B-Raf inhibitors may be determined by the following *in vitro* assays:

Fluorescence anisotropy kinase binding assay

- 10 The kinase enzyme, fluorescent ligand and a variable concentration of test compound are incubated together to reach thermodynamic equilibrium under conditions such that in the absence of test compound the fluorescent ligand is significantly (>50%) enzyme bound and in the presence of a sufficient concentration (>10x K_d) of a potent inhibitor the anisotropy of the unbound fluorescent ligand is measurably different from the bound value.

- 15 The concentration of kinase enzyme should preferably be $\geq 1 \times K_d$. The concentration of fluorescent ligand required will depend on the instrumentation used, and the fluorescent and physicochemical properties. The concentration used must be lower than the concentration of kinase enzyme, and preferably less than half the kinase enzyme concentration. A typical protocol is:

- 20 All components dissolved in Buffer of composition 50 mM HEPES, pH 7.5, 1mM CHAPS, 10 mM $MgCl_2$.

 B-Raf Enzyme concentration: 1 nM

 Fluorescent ligand concentration: 0.5 nM

 Test compound concentration: 0.1 nM - 100 μ M


- 25 Components incubated in 10 μ l final volume in LJL HE 384 type B black microtitre plate until equilibrium reached (Over 3 h, up to 30 h)

 Fluorescence anisotropy read in LJL Acquest.

Definitions: K_i = dissociation constant for inhibitor binding

K_f = dissociation constant for fluorescent ligand binding

- 30 The fluorescent ligand is the following compound:



which is derived from 5-[2-(4-aminomethylphenyl)-5-pyridin-4-yl-1H-imidazol-4-yl]-2-chlorophenol and rhodamine green.

5 Raf Kinase assay

- Activity of human recombinant B-Raf protein was assessed in vitro by assay of the incorporation of radiolabelled phosphate to recombinant MAP kinase (MEK), a known physiologic substrate of B-Raf. Catalytically active human recombinant B-Raf protein was obtained by purification from Sf9 insect cells infected with a human B-Raf recombinant baculovirus expression vector. To ensure that all substrate phosphorylation resulted from B-Raf activity, a catalytically inactive form of MEK was utilised. This protein was purified from bacterial cells expression mutant inactive MEK as a fusion protein with glutathione-S-transferase (GST-kdMEK).
- 15 **Method:** Standard assay conditions of B-Raf catalytic activity utilised 3ug of GST-kdMEK, 10uM ATP and 2uCi ^{33}P -ATP, 50mM MOPS, 0.1mM EDTA, 0.1M sucrose, 10mM MgCl_2 plus 0.1% dimethylsulphoxide (containing compound where appropriate) in a total reaction volume of 30ul. Reactions were incubated at 25°C for 90 minutes and reactions terminated by addition of EDTA to a final
- 20 concentration of 50uM. 10ul of reaction was spotted to P30 phosphocellulose paper and air dried. Following four washes in ice cold 10% trichloroacetic acid, 0.5% phosphoric acid, papers were air dried prior to addition of liquid scintillant and measurement of radioactivity in a scintillation counter.

- Results:** The compounds of the examples were found to be effective in either
- 25 of the above assays having K_d 's (in the case of the binding assay) or IC_{50} 's (in the kinase assay) of $< 3 \mu\text{M}$.

The activity of compounds as Raf inhibitors may also be determined by the assays described in WO 99/10325; McDonald, O.B., Chen, W.J., Ellis, B., Hoffman, C., Overton, L., Rink, M., Smith, A., Marshall, C.J. and Wood, E.R. (1999) A
5 scintillation proximity assay for the Raf/MEK/ERK kinase cascade: high throughput screening and identification of selective enzyme inhibitors, *Anal. Biochem.* 268: 318-329 and AACR meeting New Orleans 1998 Poster 3793.

The neuroprotective properties of B-Raf inhibitors may be determined by the
10 following *in vitro* assay:

Neuroprotective properties of B-Raf inhibitors in rat hippocampal slice cultures

Organotypic cultures provide an intermediate between dissociated neuronal cell cultures and *in-vivo* models of oxygen and glucose deprivation (OGD). The
15 majority of glial-neuronal interactions and neuronal circuitry are maintained in cultured hippocampal slices, so facilitating investigation of the patterns of death among differing cell types in a model that resembles the *in vivo* situation. These cultures allow the study of delayed cellular damage and death 24 hours, or more, post-insult and permit assessment of the consequences of long-term alterations in
20 culture conditions. A number of laboratories have reported delayed neuronal damage in response to OGD in organotypic cultures of the hippocampus (Vornov *et al.*, *Stroke*, 1994, 25, 57-465; Newell *et al.*, *Brain Res.*, 1995, 676, 38-44). Several classes of compounds have been shown to protect in this model, including EAA antagonists (Strasser *et al.*, *Brain Res.*, 1995, 687, 167-174), Na channel blockers
25 (Tasker *et al.*, *J. Neurosci.*, 1992, 12, 98-4308) and Ca channel blockers (Pringle *et al.*, *Stroke*, 1996, 7, 2124-2130). To date, relatively little is known of the roles of intracellular kinase mediated signalling pathways in neuronal cell death in this model.

Method: Organotypic hippocampal slice cultures were prepared using the
30 method of Stoppini *et al.*, *J. Neurosci. Methods*, 1995, 37, 173-182. Briefly, 400


micron sections prepared from hippocampi of 7-8 day postnatal Sprague Dawley rats are cultured on semiporous membranes for 9-12 days. OGD is then induced by incubation in serum and glucose-free medium in an anaerobic chamber for 45 minutes. Cultures are then returned to the air / CO₂ incubator for 23 hours before
5 analysis. Propidium iodide (PI) is used as an indicator of cell death. PI is non toxic to neurones and has been used in many studies to ascertain cell viability. In damaged neurons PI enters and binds to nucleic acids. Bound PI shows increased emission at 635nm when excited at 540nm. One PI fluorescence image and one
10 white light image are taken and the proportion of cell death analysed. The area of region CA1 is defined from the white light image and superimposed over the PI image. The PI signal is thresholded and area of PI damage expressed as a percentage of the CA1 area. Correlation between PI fluorescence and histologically confirmed cell death has been validated previously by Nissl-staining using cresyl fast violet (Newell *et al.*, *J. Neurosci.*, 1995, 15, 7702-7711).

15 It is to be understood that the present invention covers all combinations of particular and preferred groups described herein above.

Throughout the specification and the claims which follow, unless the context requires otherwise, the word 'comprise', and variations such as 'comprises' and 'comprising', will be understood to imply the inclusion of a stated integer or step or
20 group of integers but not to the exclusion of any other integer or step or group of integers or steps.

Claims:

1. A compound of formula (I):



(I)

wherein

X is O, CH₂, S or NH, or the moiety X-R¹ is hydrogen;

V is CH or N;

R¹ is hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aryl, arylC₁₋₆alkyl, heterocyclyl, heterocyclylC₁₋₆alkyl, heteroaryl or heteroarylC₁₋₆alkyl any of which except hydrogen may be optionally substituted;

R² and R³ independently represent hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aryl, arylC₁₋₆alkyl, heteroaryl, heteroarylC₁₋₆alkyl, heterocyclyl, or heterocyclylC₁₋₆alkyl any one of which except hydrogen may be optionally substituted, or R² and R³ together with the nitrogen atom to which they are attached form a 4- to 10-membered optionally substituted monocyclic or bicyclic ring;


Ar is an aryl or heteroaryl ring either of which may be optionally substituted;

one of X₁ and X₂ is N and the other is NR⁴, wherein R⁴ is hydrogen, C₁₋₆alkyl, or arylC₁₋₆alkyl;

or a pharmaceutically acceptable salt thereof.

2. A compound according to claim 1 wherein X-R¹ is hydrogen.

3. A compound according to any one of the preceding claims wherein Ar is optionally substituted phenyl.
4. A compound according to claim 3 wherein Ar is substituted by up to 3 substituents
5 independently selected from halo, hydroxy, hydroxy C₁₋₆alkyl, hydroxyimino C₁₋₆alkyl, and C₁₋₆alkoxy.
5. A compound as claimed in any one of the preceding claims wherein R² and R³
10 independently represent hydrogen, C₁₋₆alkyl, aryl, heteroarylC₁₋₆alkyl, any of which except hydrogen can be optionally substituted, or R² and R³ together with the nitrogen atom to which they are attached form an optionally substituted 5 or 6 membered monocyclic or bicyclic ring for example piperidine.
6. A compound as described in the Examples.
15
7. A pharmaceutical composition comprising a compound according to any one of claims 1 to 6 or a pharmaceutically acceptable salt thereof and a pharmaceutically acceptable carrier.
8. The use of a compound according to any one of claims 1 to 6 or a
20 pharmaceutically acceptable salt thereof in the manufacture of a medicament for the prophylactic or therapeutic treatment of any disease state in a human, or other mammal, which is exacerbated or caused by a neurotraumatic event.
9. The use of a compound according to any one of claims 1 to 6 or a
25 pharmaceutically acceptable salt thereof in the manufacture of a medicament for the prophylactic or therapeutic treatment of cancer.
10. A compound of formula (II):
30



(II)

wherein X , V , R^1 , Ar , X_1 and X_2 are as defined for formula (I) and R is hydrogen, C_{1-6} alkyl or aryl C_{1-6} alkyl.

INTERNATIONAL SEARCH REPORT

Int'l Application No
PCT/GB 01/00916

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 C07D401/04 C07D401/14 A61K31/44 A61P35/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BEILSTEIN Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 12876 A (MERCK & CO INC ;SELNICK HAROLD G (US); CLAREMON DAVID A (US); LIVE) 10 April 1997 (1997-04-10) page 1 -page 3; claim 21	1-10
A	WO 99 32436 A (BAYER AG) 1 July 1999 (1999-07-01) page 1 -page 2; claim 1	1-10
A	GB 2 306 108 A (MERCK & CO INC) 30 April 1997 (1997-04-30) page 1 -page 2; claim 1	1-10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *Z* document member of the same patent family

Date of the actual completion of the international search

16 May 2001

Date of mailing of the international search report

25/05/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Seelmann, I

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No
PCT/GB 01/00916

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9712876 A	10-04-1997	AU 702146 B	11-02-1999
		AU 7514396 A	28-04-1997
		CA 2234066 A	10-04-1997
		CN 1203590 A	30-12-1998
		CZ 9801043 A	15-07-1998
		EP 0854870 A	29-07-1998
		JP 11514353 T	07-12-1999
		NO 981528 A	05-06-1998
		PL 326025 A	17-08-1998
		SK 43598 A	04-11-1998
		HU 9902294 A	28-04-2000
		US 6083949 A	04-07-2000
		US 5717100 A	10-02-1998
WO 9932436 A	01-07-1999	AU 1905499 A	12-07-1999
		BG 104599 A	30-03-2001
		CN 1283180 T	07-02-2001
		DE 1049664 T	03-05-2001
		EP 1049664 A	08-11-2000
		ES 2153809 T	16-03-2001
		NO 20003230 A	21-08-2000
		TR 200002616 T	21-11-2000
GB 2306108 A	30-04-1997	US 5760623 A	02-06-1998